

Title	ABOUT THE JAMES CONSTANT OF ABSOLUTE NORMED SPACES(II) (Nonlinear Analysis and Convex Analysis)
Author(s)	MITANI, KEN-ICHI; SAITO, KICHI-SUKE; SUZUKI, TOMONARI
Citation	数理解析研究所講究録 (2008), 1611: 128-133
Issue Date	2008-09
URL	http://hdl.handle.net/2433/140046
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

ABOUT THE JAMES CONSTANT OF ABSOLUTE NORMED SPACES II

KEN-ICHI MITANI, KICHI-SUKE SAITO, AND TOMONARI SUZUKI

ABSTRACT. In this note, we describe some recent results concerning James constant of absolute norms on \mathbb{R}^2 and the 2-dimensional Lorentz sequence spaces.

1. INTRODUCTION

A Banach space X is called *uniformly non-square* if there is a $\delta > 0$ such that if $x, y \in S_X$ then $\|x + y\|/2 \leq 1 - \delta$ or $\|x - y\|/2 \leq 1 - \delta$, where $S_X = \{x \in X : \|x\| = 1\}$. Gao and Lau [4] introduced the *James constant* of a Banach space X as follows:

$$J(X) = \sup \left\{ \min \{ \|x + y\|, \|x - y\| \} : x, y \in S_X \right\}.$$

We shall collect some properties about James constant:

- (1) For any Banach space X we have $\sqrt{2} \leq J(X) \leq 2$.
- (2) If X is a Hilbert space, then $J(X) = \sqrt{2}$.
- (3) $J(X) < 2$ if and only if X is uniformly non-square.
- (4) If $1 \leq p \leq \infty$ and $\dim L_p \geq 2$, then

$$J(L_p) = \max\{2^{1/p}, 2^{1/p'}\}$$

where $1/p + 1/p' = 1$.

In this note, we describe some recent results concerning the James constant of absolute norms on \mathbb{R}^2 and the 2-dimensional Lorentz sequence spaces.

A norm $\|\cdot\|$ on \mathbb{R}^2 is said to be *absolute* if $\|(x, y)\| = \|(|x|, |y|)\|$ for all $x, y \in \mathbb{R}$, and *normalized* if $\|(1, 0)\| = \|(0, 1)\| = 1$. The ℓ_p -norms $\|\cdot\|_p$ are

such examples:

$$\|(x, y)\|_p = \begin{cases} (|x|^p + |y|^p)^{1/p} & \text{if } 1 \leq p < \infty, \\ \max\{|x|, |y|\} & \text{if } p = \infty. \end{cases}$$

Let AN_2 be the family of all absolute normalized norms on \mathbb{R}^2 . Bonsall and Duncan [2] showed that for any absolute normalized norm on \mathbb{R}^2 there corresponds a continuous convex function on $[0, 1]$ with some appropriate conditions as follows. Let Ψ_2 be the family of all continuous convex functions on $[0, 1]$ such that $\psi(0) = \psi(1) = 1$ and $\max\{1 - t, t\} \leq \psi(t) \leq 1$. Then AN_2 and Ψ_2 are in a one to one correspondence under the equation

$$(1) \quad \psi(t) = \|(1 - t, t)\| \quad (0 \leq t \leq 1).$$

Indeed, for any $\|\cdot\| \in AN_2$ we put ψ as (1). Then $\psi \in \Psi_2$. Also, for all $\psi \in \Psi_2$ we define

$$\|(x, y)\|_\psi = \begin{cases} (|x| + |y|)\psi\left(\frac{|y|}{|x| + |y|}\right) & \text{if } (x, y) \neq (0, 0), \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

Then $\|\cdot\|_\psi \in AN_2$, and $\|\cdot\|_\psi$ satisfies (1). From this result, we can consider many non- ℓ_p -type norms easily. The functions which correspond with the ℓ_p -norms $\|\cdot\|_p$ on \mathbb{R}^2 are

$$\psi_p(t) = \begin{cases} \{(1 - t)^p + t^p\}^{1/p} & \text{if } 1 \leq p < \infty, \\ \max\{1 - t, t\} & \text{if } p = \infty. \end{cases}$$

For $0 < \omega < 1$ and $1 \leq q < \infty$, the 2-dimensional Lorentz sequence space $d^{(2)}(\omega, q)$ is \mathbb{R}^2 with the norm

$$\|x\|_{\omega, q} = (x_1^{*q} + \omega x_2^{*q})^{1/q}, \quad x = (x_1, x_2) \in \mathbb{R}^2,$$

where (x_1^*, x_2^*) is the nonincreasing rearrangement of $(|x_1|, |x_2|)$; that is, $x_1^* = \max\{|x_1|, |x_2|\}$ and $x_2^* = \min\{|x_1|, |x_2|\}$.

Note here that the norm $\|\cdot\|_{\omega,q}$ of $d^{(2)}(\omega, q)$ is a symmetric absolute normalized norm on \mathbb{R}^2 , and the corresponding convex function is given by

$$\psi_{\omega,q}(t) = \begin{cases} ((1-t)^q + \omega t^q)^{1/q} & \text{if } 0 \leq t \leq 1/2, \\ (t^q + \omega(1-t)^q)^{1/q} & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

2. JAMES CONSTANT OF ABSOLUTE NORMALIZED NORMS ON \mathbb{R}^2

For a norm $\|\cdot\|$ on \mathbb{R}^2 , we write $J(\|\cdot\|)$ for $J((\mathbb{R}^2, \|\cdot\|))$. Mitani and Saito [6] characterized the James constant of $(\mathbb{R}^2, \|\cdot\|_\psi)$ in terms of ψ .

Theorem 1 ([6]). *Let $\psi \in \Psi_2$. If ψ is symmetric with respect to $t = 1/2$, then*

$$J(\|\cdot\|_\psi) = \max_{0 \leq t \leq 1/2} \frac{2-2t}{\psi(t)} \psi\left(\frac{1}{2-2t}\right).$$

Example 2. Let $1 \leq p \leq \infty$ and $1/p + 1/p' = 1$. Then

$$(2) \quad J(\|\cdot\|_p) = \max\{2^{1/p}, 2^{1/p'}\}.$$

Indeed, we define a function f on $[0, 1/2]$ as follows:

$$\begin{aligned} f(t) &= \frac{2-2t}{\psi_p(t)} \psi_p\left(\frac{1}{2-2t}\right) \\ &= \left(\frac{1 + (1-2t)^p}{(1-t)^p + t^p}\right)^{1/p}. \end{aligned}$$

If $1 \leq p \leq 2$, then f is the maximum at $t = 0$ and

$$J(\|\cdot\|_{\psi_p}) = f(0) = 2^{1/p}.$$

If $p \geq 2$, then f is the maximum at $t = 1/2$ and

$$J(\|\cdot\|_{\psi_p}) = f(1/2) = 2^{1/p'}.$$

Thus we obtain (2).

Example 3. Let $1/2 \leq \lambda \leq 1$. We define a function φ_λ as

$$\varphi_\lambda(t) = \max\{1 - t, t, \lambda\}.$$

Then it is obvious that $\varphi_\lambda \in \Psi_2$. The corresponding absolute normalized norm $\|\cdot\|_{\varphi_\lambda}$ is

$$\|\cdot\|_{\varphi_\lambda} = \max\{\|\cdot\|_\infty, \lambda\|\cdot\|_1\}.$$

Then

$$J(\|\cdot\|_{\varphi_\lambda}) = \begin{cases} 1/\lambda & \text{if } 1/2 \leq \lambda \leq 1/\sqrt{2}, \\ 2\lambda & \text{if } 1/\sqrt{2} \leq \lambda \leq 1. \end{cases}$$

3. JAMES CONSTANT OF 2-DIMENSIONAL LORENTZ SEQUENCE SPACES

Kato and Maligranda [5] calculated $d^{(2)}(\omega, q)$ in the case where $q \geq 2$, that is, they proved that if $0 < \omega < 1$ and $q \geq 2$, then

$$J(d^{(2)}(\omega, q)) = 2 \left(\frac{1}{1 + \omega} \right)^{1/q}.$$

However, from Theorem 1 we obtain the following.

Lemma 4. For $0 < \omega < 1$ and $1 \leq q < \infty$,

$$J(d^{(2)}(\omega, q)) (= J(\|\cdot\|_{\psi_{\omega,q}})) = \max_{0 \leq t \leq 1/2} \frac{2 - 2t}{\psi_{\omega,q}(t)} \psi_{\omega,q} \left(\frac{1}{2 - 2t} \right)$$

holds.

By using this lemma, we calculate $J(d^{(2)}(\omega, q))$ in the case where $1 \leq q < 2$.

Theorem 5 ([9], cf. [6, 12]). Let $1 \leq q < 2$. (i) If $0 < \omega \leq (\sqrt{2} - 1)^{2-q}$, then

$$J(d^{(2)}(\omega, q)) = 2 \left(\frac{1}{1 + \omega} \right)^{1/q}.$$

(ii) If $(\sqrt{2} - 1)^{2-q} < \omega < 1$, then there exists a unique solution s_0 of the equation


$$(1 + s_0)^{q-1}(1 - \omega s_0^{q-1}) = \omega(1 - s_0)^{q-1}(1 + \omega s_0^{q-1}), \quad 0 < s_0 < \omega^{1/(2-q)}.$$

(ii-a) If $(\sqrt{2} - 1)^{2-q} < \omega \leq \sqrt{2}^q - 1$, then

$$J(d^{(2)}(\omega, q)) = \max \left\{ \left(\frac{2(1 + s_0)^{q-1}}{1 + \omega s_0^{q-1}} \right)^{1/q}, 2 \left(\frac{1}{1 + \omega} \right)^{1/q} \right\}.$$

(ii-b) If $\sqrt{2}^q - 1 < \omega < 1$, then

$$J(d^{(2)}(\omega, q)) = \left(\frac{2(1 + s_0)^{q-1}}{1 + \omega s_0^{q-1}} \right)^{1/q}.$$



REFERENCES

- [1] B. Beauzamy, *Introduction to Banach Spaces and Their Geometry*, 2nd ed., North-Holland, Amsterdam-New York-Oxford, 1985.
- [2] F. F. Bonsall and J. Duncan, *Numerical Ranges II*, London Math. Soc. Lecture Note Series, Vol. 10, 1973.
- [3] E. Casini, *About some parameters of normed linear spaces*, Atti Accad. Naz. Lincei, VIII. Ser., Rend., Cl. Sci. Fis. Mat. Nat., **80**(1986), 11–15.
- [4] J. Gao and K. S. Lau, *On the geometry of spheres in normed linear spaces*, J. Austral. Math. Soc., A **48**(1990), 101–112.
- [5] M. Kato and L. Maligranda, *On James and Jordan-von Neumann constants of Lorentz sequence spaces*, J. Math. Anal. Appl., **258**(2001), 457–465.
- [6] K.-I. Mitani and K.-S. Saito, *The James constant of absolute norms on \mathbb{R}^2* , J. Nonlinear Convex Anal., **4**(2003), 399–410.
- [7] K.-I. Mitani and K.-S. Saito, *About the James constant of absolute normed spaces*, RIMS Kokyuroku, **1365**(2004), 33–40.

- [8] K.-I. Mitani and K.-S. Saito, *The James constant and von Neumann-Jordan constant of absolute norms on \mathbb{R}^2* , Proceedings of the International Symposium on Banach and Function Spaces, 315–321, Yokohama Publ., Yokohama, 2004.
- [9] K.-I. Mitani, K.-S. Saito and T. Suzuki, *On the calculation of the James constant of Lorentz sequence spaces*, submitted.
- [10] K.-S. Saito, M. Kato and Y. Takahashi, *Von Neumann-Jordan Constant of absolute normalized norms on \mathbb{C}^2* , J. Math. Anal. Appl., **244**(2000), 515–532.
- [11] K.-S. Saito, M. Kato and Y. Takahashi, *Absolute norms on \mathbb{C}^n* , J. Math. Anal. Appl., **252**(2000), 879–905.
- [12] T. Suzuki, A. Yamano and M. Kato, *The James constant of 2-dimensional Lorentz sequence spaces*, Bull. Kyushu Inst. Technol. Pure Appl. Math., **53**(2006), 15–24.

(K.-I. Mitani) NIIGATA INSTITUTE OF TECHNOLOGY, KASHIWAZAKI, NIIGATA 945-1195, JAPAN

E-mail address: mitani@adm.niit.ac.jp

(K.-S. Saito) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, NIIGATA UNIVERSITY, NIIGATA 950-2181, JAPAN

E-mail address: saito@math.sc.niigata-u.ac.jp

(T. Suzuki) DEPARTMENT OF MATHEMATICS, KYUSHU INSTITUTE OF TECHNOLOGY, KITAKYUSHU 804-8550, JAPAN

E-mail address: suzuki-t@mns.kyutech.ac.jp